Journal of MOLECULAR BIOLOGY

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
C. R. Cantor
P. Chambon
W. W. Franke
M. Gellert
G. A. Gilbert
M. Gottesman
I. Herskowitz
R. Huber
H. E. Huxley
A. Klug
V. Luzzati
J. H. Miller
M. F. Moody
K. Simons

H
01
T73

VOLUME 169

5 September to 5 October 1983

ACADEMIC PRESS
London New York San Francisco

Copyright © 1983, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 169

Number 1, 5 September

Trifonov, E. N. and Bolshoi, G., Open and Closed 5 S Ribosomal RNA, the Only Two Universal Structures Encoded in the Nucleotide Sequences 1

McLachlan, A. D., Analysis of Gene Duplication Repeats in the Myosin Rod 15

Shapiro, S. G., Schon, E. A., Townes, T. M. and Lingrel, J. B., Sequence and Linkage of the Goat ϵ^{I} and ϵ^{II} β-Globin Genes 31

Hinton, D. M. and Musso, R. E., Specific *in Vitro* Transcription of the Interaction Sequence IS2 53

Appendix 69

Hall, L. M. C., Mason, P. J. and Spierer, P., Transcripts, Genes and Bands in 315,000 Base-pairs of *Drosophila* DNA 83

Heuser, J. E. and Cooke, R., Actin–Myosin Interactions Visualized by the Quick-freeze, Deep-etch Replica Technique 97

Heuser, J. E., Structure of the Myosin Crossbridge Lattice in Insect Flight Muscle 123

Heuser, J. E., Procedure for Freeze-drying Molecules Adsorbed to Mica Flakes 155

Peeters, B. P. H., Konings, R. N. H. and Schoenmakers, J. G. G., Characterization of the DNA Binding Protein Encoded by the N-specific Filamentous *Escherichia coli* Phage IKe. Binding Properties of the Protein and Nucleotide Sequence of the Gene 197

Bartenev, V. N., Golovanov, E. I., Kapitonova, K. A., Mokulskii, M. A., Volkova, L. I. and Skuratovskii, I. Ya., Structure of the *B* DNA Cationic Shell as Revealed by an X-ray Diffraction Study of CsDNA. Sequence-specific Cationic Stabilization of *B* Form DNA. . 217

Cohen, H. A., Chiu, W. and Hosoda, J., Structural Analysis of T4 DNA Helix Destabilizing Protein (gp32*I) Crystal by Electron Microscopy . 235

Douthwaite, S., Christensen, A. and Garrett, R. A., Higher Order Structure in the 3′-Minor Domain of Small Subunit Ribosomal RNAs from a Gram Negative Bacterium, a Gram Positive Bacterium and a Eukaryote 249

KUWAJIMA, K. and BALDWIN, R. L., Nature and Locations of the Most Slowly Exchanging Peptide NH Protons in Residues 1 to 19 of Ribonuclease S 281

KUWAJIMA, K. and BALDWIN, R. L., Exchange Behaviour of the H-bonded Amide Protons in the 3 to 13 Helix of Ribonuclease S 299

ENGLANDER, J. J., ROGERO, J. R. and ENGLANDER, S. W., Identification of an Allosterically Sensitive Unfolding Unit in Hemoglobin 325

LETTERS TO THE EDITOR

KISELEV, N. A., ORLOVA, E. V., STEL'MASCHUK, V. YA., VASILIEV, V. D., SELIVANOVA, O. M., KOSYKH, V. P., PUSTOVSKIKH, A. I. and KIRICHUK, V. S., Computer Averaging of 50 S Ribosomal Subunit Electron Micrographs 345

BOLOGNESI, M., GATTI, G., CODA, A., AVIGLIANO, L., MARCOZZI, G. and FINAZZI-AGRÒ, A., A New Crystalline Modification of the Copper Enzyme Ascorbate Oxidase 351

NUMBER 2, 15 SEPTEMBER 1983

AUSTIN, S. and ABELES, A., Partition of Unit-copy Miniplasmids to Daughter Cells. I. P1 and F Miniplasmids Contain Discrete, Interchangeable Sequences Sufficient to Promote Equipartition . . 353

AUSTIN, S. and ABELES, A., Partition of Unit-copy Miniplasmids to Daughter Cells. II. The Partition Region of Miniplasmid P1 Encodes an Essential Protein and a Centromere-like Site at which it Acts . . 373

HORIUCHI, K., Co-evolution of a Filamentous Bacteriophage and its Defective Interfering Particles 389

SERDYUK, I. N., AGALAROV, S. CH., SEDELNIKOVA, S. E., SPIRIN, A. S. and MAY, R. P., Shape and Compactness of the Isolated Ribosomal 16 S RNA and its Complexes with Ribosomal Proteins 409

URIG, M. A., BROWN, S. M., TEDESCO, P. and WOOD, W. B., Attachment of Tail Fibers in Bacteriophage T4 Assembly. Identification of the Baseplate Protein to which Tail Fibers Attach 427

FREUNDLICH, A. and SQUIRE, J. M., Three-dimensional Structure of the Insect (*Lethocerus*) Flight Muscle M-Band 439

APPLEGATE, D. and REISLER, E., Crossbridge Release and α-Helix–Coil Transition in Myosin and Rod Minifilaments 455

HUXLEY, H. E., SIMMONS, R. M., FARUQI, A. R., KRESS, M., BORDAS, J. and KOCH, M. H. J., Changes in the X-ray Reflections from Contracting Muscle During Rapid Mechanical Transients and their Structural Implications 469

MANOIL, C., A Genetic Approach to Defining the Sites of Interaction of a Membrane Protein with Different External Agents. 507

GUSS, J. M. and FREEMAN, H. C., Structure of Oxidized Poplar Plastocyanin at 1·6 Å Resolution. 521

BRAYER, G. D. and MCPHERSON, A., Refined Structure of the Gene *5* DNA Binding Protein from Bacteriophage fd 565

BARDSLEY, W. G., SOLANO-MUÑOZ, F., WRIGHT, A. J. and MCGINLAY, P. B., Inhibition of Enzyme-catalysed Reactions by Excess Substrate. A Theoretical and Monte Carlo Study of Turning Points in $v(S)$ Graphs . 597

Appendix 611

KREBS, H., SCHMID, F. X. and JAENICKE, R., Folding of Homologous Proteins. The Refolding of Different Ribonucleases is Independent of Sequence Variations, Proline Content and Glycosylation 619

LETTERS TO THE EDITOR

AMIT, A. G., MARIUZZA, R. A., SALUDJIAN, P., POLJAK, R. J., LAMOYI, E. and NISONOFF, A., Preliminary Crystallographic Study of the Fab Fragment of a Monoclonal Anti-phenylarsonate Antibody Possessing a Major Idiotypic Specificity 637

ERRATUM 639

NUMBER 3, 25 SEPTEMBER 1983

SMITH, M. M. and MURRAY, K., Yeast H3 and H4 Histone Messenger RNAs are Transcribed from Two Non-allelic Gene Sets 641

SMITH, M. M. and ANDRÉSSON, O. S., DNA Sequences of Yeast H3 and H4 Histone Genes from Two Non-allelic Gene Sets Encode Identical H3 and H4 Proteins 663

ALONSO, A., JORCANO, J. L., BECK, E. and SPIESS, E., Isolation and Characterization of *Drosophila melanogaster* U2 Small Nuclear RNA Genes 691

HILLEN, W., GATZ, C., ALTSCHMIED, L., SCHOLLMEIER, K. and MEIER, I., Control of Expression of the Tn*10*-encoded Tetracycline Resistance Genes. Equilibrium and Kinetic Investigation of the Regulatory Reactions 707

FLICKER, P. F., WALLIMANN, T. and VIBERT, P., Electron Microscopy of Scallop Myosin. Location of Regulatory Light Chains 723

BORKAKOTI, N., PALMER, R. A., HANEEF, I. and MOSS, D. S., Specificity of Pancreatic Ribonuclease-A. An X-ray Study of a Protein–Nucleotide Complex. 743

Ohlendorf, D. H., Anderson, W. F., Lewis, M., Pabo, C. O. and Matthews, B. W., Comparison of the Structures of Cro and λ Repressor Proteins from Bacteriophage λ 757

Letter to the Editor

Wrigley, N. G., Brown, E. B. and Skehel, J. J., Electron Microscopic Evidence for the Axial Rotation and Inter-domain Flexibility of the Fab Regions of Immunoglobulin G 771

Number 4, 5 October 1983

Horowitz, H., Van Arsdell, J. and Platt, T., Nucleotide Sequence of the *trpD* and *trpC* Genes of *Salmonella typhimurium* 775

Spena, A., Viotti, A. and Pirrotta, V., Two Adjacent Genomic Zein Sequences: Structure, Organization and Tissue-specific Restriction Pattern 799

Mitra, A. K., Arnott, S. and Sheehan, J. K., Hyaluronic Acid: Molecular Conformation and Interactions in the Tetragonal Form of the Potassium Salt-containing Extended Chains 813

Mitra, A. K., Raghunathan, S., Sheehan, J. K. and Arnott, S., Hyaluronic Acid: Molecular Conformations and Interactions in the Orthorhombic and Tetragonal Forms Containing Sinuous Chains . . 829

Arnott, S., Mitra, A. K. and Raghunathan, S., Hyaluronic Acid Double Helix 861

Mitra, A. K., Arnott, S., Atkins, E. D. T. and Isaac, D. H., Dermatan Sulfate: Molecular Conformations and Interactions in the Condensed State 873

Millane, R. P., Mitra, A. K. and Arnott, S., Chondroitin 4-Sulfate: Comparison of the Structures of the Potassium and Sodium Salts . . 903

Braun, W., Wider, G., Lee, K. H. and Wüthrich, K., Conformation of Glucagon in a Lipid–Water Interphase by ^{1}H Nuclear Magnetic Resonance 921

Wüthrich, K., Billeter, M. and Braun, W., Pseudo-structures for the 20 Common Amino Acids for Use in Studies of Protein Conformations by Measurements of Intramolecular Proton–Proton Distance Constraints with Nuclear Magnetic Resonance 949

Griffith, J. P., Lee, B., Murdock, A. L. and Amelunxen, R. E., Molecular Symmetry of Glyceraldehyde-3-phosphate Dehydrogenase from *Bacillus coagulans* 963

Roberts, M. P., Falvey, E., Calvet, J. P. and Pederson, T., A Sequence Related to 4·5 S RNA and the B1 Family of Repeated DNA in the 5′ Flanking Region of the Mouse β-Globin Gene. 975

Index to Volume 169

Abeles, A. (see also Austin, S.), 353, 373
Actin–myosin interactions visualized by the quick-freeze, deep-etch replica technique, 97
Agalarov, S. Ch. (see also Serdyuk, I. N.), 409
Allosterically sensitive unfolding unit in hemoglobin, identification, 325
Alonso, A., Jorcano, J. L., Beck, E. & Spiess, E., Isolation and characterization of *Drosophila melanogaster* U2 small nuclear RNA genes, 691
Altschmied, L. (see also Hillen, W.), 707
Amelunxen, R. E. (see also Griffith, J. P.), 963
Amide protons, H-bonded, in the 3 to 13 helix of ribonuclease S, exchange behaviour, 299
Amino acid pseudo-structures for use in studies of protein conformations by measurements of intramolecular proton–proton distance constraints with n.m.r., 949
Amit, A. G., Mariuzza, R. A., Saludjian, P., Poljak, R. J., Lamoyi, E. & Nisonoff, A., Preliminary crystallographic study of the Fab fragment of a monoclonal anti-phenylarsonate antibody possessing a major idiotypic specificity, 637
Anderson, W. F. (see also Ohlendorf, D. H.), 757
Andrésson, O. S. (see also Smith, M. M.), 663
Anti-phenylarsonate antibody, monoclonal, preliminary crystallographic study of the Fab fragment possessing a major idiotypic specificity, 637
Applegate, D. & Reisler, E., Crossbridge release and α-helix–coil transition in myosin and rod minifilaments, 455
Arnott, S., Mitra, A. K. & Raghunathan, S., Hyaluronic acid double helix, 861
— (see also Millane, R. P.), 903
— (see also Mitra, A. K.), 813, 829, 873
Ascorbate oxidase, a new crystalline modification, 351
Atkins, E. D. T. (see also Mitra, A. K.), 873
Attachment of tail fibers in bacteriophage T4 assembly. Identification of the baseplate protein to which tail fibers attach, 427
Austin, S. & Abeles, A., Partition of unit-copy miniplasmids to daughter cells. I. P1 and F miniplasmids contain discrete, interchangeable sequences sufficient to promote equipartition, 353
— & —, — — —-— — — — —. II. The partition region of miniplasmid P1 encodes an essential protein and a centromere-like site at which it acts, 373
Avigliano, L. (see also Bolognesi, M.), 351

B DNA cationic shell, structure as revealed by X-ray diffraction study of CsDNA, 217
Bacillus coagulans, molecular symmetry of glyceraldehyde-3-phosphate dehydrogenase, 963
Bacteriophage fd, refined structure of the gene *5* DNA binding proteins, 565
—, filamentous, and its defective interfering particles, co-evolution, 389
— λ, comparison of the structures of cro and λ repressor proteins, 757
— T4 assembly, identification of the baseplate protein to which tail fibers attach, 427
Baldwin, R. L. (see also Kuwajima, K.), 281, 299
Bands in 315,000 base-pairs of *Drosophila* DNA, 83
Bardsley, W. G., Solano-Muñoz, F., Wright, A. J. & McGinlay, P. B., Inhibition of enzyme-catalysed reactions by excess substrate. A theoretical and Monte Carlo study of turning points in $v(S)$ graphs, 597
Bartenev, V. N., Golovanov, E. I., Kapitonova, K. A., Mokulskii, M. A., Volkova, L. I. & Skuratovskii, I. Ya., Structure of the *B* DNA cationic shell as revealed by an X-ray diffraction study of CsDNA. Sequence-specific cationic stabilization of *B* form DNA, 217
Baseplate protein to which bacteriophage T4 tail fibers attach in assembly, 427
Beck, E. (see also Alonso, A.), 691
Billeter, M. (see also Wüthrich, K.), 949
Binding protein, DNA, encoded by the N-specific filamentous *Escherichia coli* phage IKe, characterization, 197
Binding protein, DNA, gene *5*, from bacteriophage fd; refined structure, 565
Bolognesi, M., Gatti, G., Coda, A., Avigliano, L., Marcozzi, G. & Finazzi-Agrò, A., A new crystalline modification of the copper enzyme ascorbate oxidase, 351
Bolshoi, G. (see also Trifonov, E. N.), 1

Bordas, J. (see also Huxley, H. E.), 469
Borkakoti, N., Palmer, R. A., Haneef, I. & Moss, D. S., Specificity of pancreatic ribonuclease-A. An X-ray study of protein–nucleotide complex, 743
Braun, W., Wider, G., Lee, K. H. & Wüthrich, K., Conformation of glucagon in a lipid–water interphase by ^{1}H nuclear magnetic resonance, 921
Braun, W. (see also Wüthrich, K.), 949
Brayer, G. D. & McPherson, A., Refined structure of the gene *5* DNA binding protein from bacteriophage fd, 565
Brown, E. B. (see also Wrigley, N. G.), 771
Brown, S. M. (see also Urig, M. A.), 427

Calvet, J. P. (see also Roberts, M. P.), 975
Cationic stabilization, sequence-specific, of *B* form DNA; structure of the *B* DNA cationic shell as revealed by X-ray diffraction of CsDNA, 217
Chiu, W. (see also Cohen, H. A.), 235
Chondroitin 4-sulfate: comparison of the structures of the potassium and sodium salts, 903
Christensen, A. (see also Douthwaite, S.), 249
Coda, A. (see also Bolognesi, M.), 351
Cohen, H. A., Chiu, W. & Hosoda, J., Structural analysis of T4 DNA helix destabilizing protein (gp32*I) crystal by electron microscopy, 235
Computer averaging of 50 S ribosomal subunit electron micrographs, 345
Conformation of glucagon in a lipid–water interphase by ^{1}H nuclear magnetic resonance, 921
—, molecular, and interactions of dermatan sulfate in the condensed state, 873
—, —, — — — hyaluronic acid in the orthorhombic and tetragonal forms containing sinuous chains, 829
—, —, — — — — — — — tetragonal form of the potassium salt containing extended chains, 813
— of proteins by n.m.r. measurements of intramolecular proton–proton distance constraints, use of pseudo-structures for the 20 common amino acids, 949
Cooke, R. (see also Heuser, J. E.), 97
Copper enzyme ascorbate oxidase, a new crystalline modification, 351
Cro and λ repressor proteins from bacteriophage λ, comparison of structures, 757
Crossbridge myosin lattice in insect flight muscle, study of structure, 123
— release and α-helix–coil transition in myosin and rod minifilaments, 455
Crystalline modification, new, of the copper enzyme ascorbate oxidase, 351
Crystallographic study, preliminary, of the Fab fragment of a monoclonal anti-phenylarsonate antibody possessing a major idiotypic specificity, 637
CsDNA, X-ray diffraction study reveals the structure of *B* DNA cationic shell; sequence-specific cationic stabilization of *B* form DNA, 217

Defective interfering particles, and filamentous bacteriophage, co-evolution, 389
DNA, *B*, structure of the cationic shell as revealed by X-ray diffraction study of CsDNA, 217
DNA binding protein encoded by the N-specific filamentous *Escherichia coli* phage IKe, characterization; binding properties of the protein and nucleotide sequence of the gene, 197
— binding protein, gene *5*, from bacteriophage fd, refined structure, 565
—, *Drosophila*, transcripts, genes and bands in 315,000 base-pairs, 83
— helix destabilizing protein (gp32*I) crystal, structural analysis by electron microscopy, 235
—, repeated, B1 family in the 5′ flanking region of the mouse β-globin gene, a sequence related to 4·5 S RNA and, 975
— sequences of yeast H3 and H4 histone genes from two non-allelic gene sets encode identical H3 and H4 proteins, 663
Dermatan sulfate: molecular conformations and interactions in the condensed state, 873
Douthwaite, S., Christensen, A. & Garrett, R. A., Higher order structure in the 3′-minor domain of small subunit ribosomal RNAs from a gram negative bacterium, a gram positive bacterium and a eukaryote, 249
Drosophila DNA, transcripts, genes and bands in 315,000 base-pairs, 83
Drosophila melanogaster U2 small nuclear RNA genes, isolation and characterization, 691

Electron micrographs of 50 S ribosomal subunit, computer averaging, 345
— microscopic evidence for the axial rotation and inter-domain flexibility of the Fab regions of immunoglobulin G, 771
— microscopy of scallop myosin; location of regulatory light chains, 723

— —, use in structural analysis of T4 DNA helix destabilizing protein (gp32*I) crystal, 235
ENGLANDER, J. J., ROGERO, J. R. & ENGLANDER, S. W., Identification of an allosterically sensitive unfolding unit in hemoglobin, 325
ENGLANDER, S. W. (see also ENGLANDER, J. J.), 325
Enzyme-catalysed reactions, inhibition by excess substrate; a theoretical and Monte Carlo study of turning points in $v(S)$ graphs, 597
ϵ^{I} and ϵ^{II} β-globin genes, goat, sequence and linkage, 31
Escherichia coli phage IKe, filamentous, N-specific, characterization of the DNA binding protein, 197
Evolution of a filamentous bacteriophage and its defective interfering particles, 389
Exchange behaviour of the H-bonded amide protons in the 3 to 13 helix of ribonuclease S, 299
Expression control of the Tn*10*-encoded tetracycline resistance genes; equilibrium and kinetic investigation of the regulatory reactions, 707

Fab regions of immunoglobin G, electron microscopic evidence for axial rotation and inter-domain flexibility, 771
— fragment of a monoclonal anti-phenylarsonate antibody possessing a major idiotypic specificity, preliminary crystallographic study, 637
FALVEY, E. (see also ROBERTS, M. P.), 975
FARUQI, A. R. (see also HUXLEY, H. E.), 469
fd (see also Bacteriophage), 565
Filamentous bacteriophage and its defective interfering particles, co-evolution, 389
— *Escherichia coli* phage IKe, characterization of the encoded DNA binding protein, 197
FINAZZI-AGRÒ, A. (see also BOLOGNESI, M.), 351
FLICKER, P. F., WALLIMANN, T. & VIBERT, P., Electron microscopy of scallop myosin. Location of regulatory light chains, 723
Folding of homologous proteins. The refolding of different ribonucleases is independent of sequence variations, proline content and glycosylation, 619
FREEMAN, H. C. (see also GUSS, J. M.), 521
Freeze-drying procedure for molecules adsorbed to mica flakes, 155
Freeze-etch replica technique for visualization of actin–myosin interactions, 97
FREUNDLICH, A. & SQUIRE, J. M., Three-dimensional structure of the insect (*Lethocerus*) flight muscle M-band, 439

GARRETT, R. A. (see also DOUTHWAITE, S.), 249
GATTI, G. (see also BOLOGNESI, M.), 351
GATZ, C. (see also HILLEN, W.), 707
Gene, β-globin, mouse, a sequence related to 4·5 S RNA and the B1 family of repeated DNA in the 5′ flanking region, 975
— duplication repeats in the myosin rod, 15
— *5* DNA binding protein from bacteriophage fd, refined structure, 565
— specifying the IKe DNA binding protein, nucleotide sequence, 197
Genes, β-globin, ϵ^{I} and ϵ^{II}, goat, sequence and linkage, 31
— coding for H3 and H4 histones of *Saccharomyces cerevisiae*, 641
— encoding for H3 and H4 histones of *Saccharomyces cerevisiae*, DNA sequences from two non-allelic gene sets encode identical H3 and H4 proteins, 663
—, tetracycline resistance, Tn*10*-encoded, control of expression; equilibrium and kinetic investigation of the regulatory reactions, 707
—, transcripts and bands in 315,000 base-pairs of *Drosophila* DNA, 83
—, *trpD* and *trpC*, of *Salmonella typhimurium*, nucleotide sequences, 775
—, U2 small nuclear RNA, *Drosophila melanogaster*, isolation and characterization, 691
Genetic approach to defining the sites of interaction of a membrane protein with different external agents, 507
Genomic zein sequences, adjacent: structure, organization and tissue-specific restriction pattern, 799
β-Globin gene, mouse, a sequence related to 4·5 S RNA and the B1 family of repeated DNA in the 5′ flanking region, 975
— genes, ϵ^{I} and ϵ^{II}, goat, sequence and linkage, 31
Glucagon in a lipid–water interphase, conformation by ^{1}H n.m.r., 921
Glyceraldehyde-3-phosphate dehydrogenase from *Bacillus coagulans*, molecular symmetry, 963
Glycosylation of ribonucleases does not affect refolding, 619
Goat ϵ^{I} and ϵ^{II} β-globin genes, sequence and linkage, 31
GOLOVANOV, E. I. (see also BARTENEV, V. N.), 217

gp32*I crystal, structural analysis by electron microscopy, 235
Gram negative bacterium, gram positive bacterium and a eukaryote, higher order structure in the 3′-minor domain of their small subunit ribosomal RNAs, 249
Griffith, J. P., Lee, B., Murdock, A. L. & Amelunxen, R. E., Molecular symmetry of glyceraldehyde-3-phosphate dehydrogenase from *Bacillus coagulans*, 963
Guss, J. M. & Freeman, H. C., Structure of oxidized poplar plastocyanin at 1·6 Å resolution, 521

^{1}H nuclear magnetic resonance determination of the conformation of glucagon in a lipid–water interphase, 921
H3 and H4 histone genes, DNA sequences from two non-allelic gene sets encode identical H3 and H4 proteins, 663
H3 and H4 histone mRNAs, yeast, are transcribed from two non-allelic gene sets, 641
Hall, L. M. C., Mason, P. J. & Spierer, P., Transcripts, genes and bands in 315,000 base-pairs of *Drosophila* DNA, 83
Haneef, I. (see also Borkakoti, N.), 743
Helix destabilizing protein (gp32*I) crystal, structural analysis by electron microscopy, 235
—, double, hyaluronic acid, 861
α-Helix–coil transition and crossbridge release in myosin and rod minifilaments, 455
Hemoglobin, identification of an allosterically sensitive unfolding unit, 325
Heuser, J. E., Procedure for freeze-drying molecules adsorbed to mica flakes, 155
—, Structure of the myosin crossbridge lattice in insect flight muscle, 123
— & Cooke, R., Actin–myosin interactions visualized by the quick-freeze, deep-etch replica technique, 97
Hillen, W., Gatz, C., Altschmied, L., Schollmeier, K. & Meier, I., Control of expression of the Tn*10*-encoded tetracycline resistance genes. Equilibrium and kinetic investigation of the regulatory reactions, 707
Hinton, D. M. & Musso, R. E., Specific *in vitro* transcription of the interaction sequence IS2, 53
Horiuchi, K., Co-evolution of a filamentous bacteriophage and its defective interfering particles, 389
Horowitz, H., Van Arsdell, J. & Platt, T., Nucleotide sequence of the *trpD* and *trpC* genes of *Salmonella typhimurium*, 775
Hosoda, J. (see also Cohen, H. A.), 235
Huxley, H. E., Simmons, R. M., Faruqi, A. R., Kress, M., Bordas, J. & Koch, M. H. J., Changes in the X-ray reflections from contracting muscle during rapid mechanical transients and their structural implications, 469
Hyaluronic acid double helix, 861
— —: molecular conformation and interactions in the tetragonal form of the potassium salt-containing extended chains, 813
— —: — conformations and interactions in the orthorhombic and tetragonal forms containing sinuous chains, 829

IKe (see also Phage), 197
Immunoglobulin G, electron microscopic evidence for the axial rotation and inter-domain flexibility of the Fab regions, 771
Insect flight muscle, structure of the myosin crossbridge lattice, 123
Insect (*Lethocerus*) flight muscle M-band, three-dimensional structure, 439
Insertion sequence IS2, specific *in vitro* transcription, 53
IS2 insertion sequence, specific *in vitro* transcription, 53
Isaac, D. H. (see also Mitra, A. K.), 873

Jaenicke, R. (see also Krebs, H.), 619
Jorcano, J. L. (see also Alonso, A.), 691

Kapitonova, K. A. (see also Bartenev, V. N.), 217
Kirichuk, V. S. (see also Kiselev, N. A.), 345
Kiselev, N. A., Orlova, E. V., Stel'maschuk, V. Ya., Vasiliev, V. D., Selivanova, O. M., Kosykh, V. P., Pustovskikh, A. I. & Kirichuk, V. S., Computer averaging of 50 S ribosomal subunit electron micrographs, 345
Koch, M. H. J. (see also Huxley, H. E.), 469
Konings, R. N. H. (see also Peeters, B. P. H.), 197
Kosykh, V. P. (see also Kiselev, N. A.), 345

Krebs, H., Schmid, F. X. & Jaenicke, R., Folding of homologous proteins. The refolding of different ribonucleases is independent of sequence variations, proline content and glycosylation, 619
Kress, M. (see also Huxley, H. E.), 469
Kuwajima, K. & Baldwin, R. L., Exchange behaviour of the H-bonded amide protons in the 3 to 13 helix of ribonuclease S, 299
Kuwajima, K. & Baldwin, R. L., Nature and locations of the most slowly exchanging peptide NH protons in residues 1 to 19 of ribonuclease S, 281

λ (see also Bacteriophage), 757
Lamoyi, E. (see also Amit, A. G.), 637
Lee, B. (see also Griffith, J. P.), 963
Lee, K. H. (see also Braun, W.), 921
Lethocerus flight muscle M-band, three-dimensional structure, 439
Lewis, M. (see also Ohlendorf, D. H.), 757
Light chain location for scallop myosin by electron microscopy, 723
Lingrel, J. B. (see also Shapiro, S. G.), 31
Lipid–water interphase glucagon, conformation by ^{1}H n.m.r., 921

McGinlay, P. B. (see also Bardsley, W. G.), 597
McLachlan, A. D., Analysis of gene duplication repeats in the myosin rod, 15
McPherson, A. (see also Brayer, G. D.), 565
Manoil, C., A genetic approach to defining the sites of interaction of a membrane protein with different external agents, 507
Marcozzi, G. (see also Bolognesi, M.), 351
Mariuzza, R. A. (see also AMIT, A. G.), 637
Mason, P. J. (see also Hall, L. M. C.), 83
Matthews, B. W. (see also Ohlendorf, D. H.), 757
May, R. P. (see also Serdyuk, I. N.), 409
M-band three-dimensional structure of insect (*Lethocerus*) flight muscle, 439
Meier, I. (see also Hillen, W.), 707
Membrane protein, a genetic approach to defining sites of interaction with different external agents, 507
Messenger RNAs, H3 and H4 histone, yeast, are transcribed from two non-allelic gene sets, 641
Mica flakes, procedure for freeze-drying adsorbed molecules, 155
Millane, R. P., Mitra, A. K. & Arnott, S., Chondroitin 4-sulfate: comparison of the structures of the potassium and sodium salts, 903
Miniplasmids, unit-copy, partition to daughter cells. P1 and F miniplasmids contain discrete, interchangeable sequences sufficient to promote equipartition, 353
—, —, — — — —. The partition region of miniplasmid P1 encodes an essential protein and a centromere-like site at which it acts, 373
Mitra, A. K., Arnott, S. & Sheehan, J. K., Hyaluronic acid: molecular conformation and interactions in the tetragonal form of the potassium salt-containing extended chains, 813
—, —, Atkins, E. D. T. & Isaac, D. H., Dermatan sulfate: molecular conformations and interactions in the condensed state, 873
—, Raghunathan, S., Sheehan, J. K. & Arnott, S., Hyaluronic acid: molecular conformations and interactions in the orthorhombic and tetragonal forms containing sinuous chains, 829
— (see also Millane, R. P.), 903
Mokulskii, M. A. (see also Bartenev, V. N.), 217
Monoclonal anti-phenylarsonate antibody, preliminary crystallographic study of the Fab fragment possessing a major idiotypic specificity, 637
Monte Carlo study of turning points in $v(S)$ graphs; inhibition of enzyme-catalysed reactions by excess substrate, 597
Moss, D. S. (see also Borkakoti, N.), 743
Mouse β-globin gene, a sequence related to 4·5 S RNA and the B1 family of repeated DNA in the 5′ flanking region, 975
Murdock, A. L. (see also Griffith, J. P.), 963
Murray, K. (see also Smith, M. M.), 641
Muscle, contracting, changes in the X-ray reflections during rapid mechanical transients and their structural implications, 469
—, insect flight, structure of the myosin crossbridge lattice, 123
—, — (*Lethocerus*) flight, M-band three-dimensional structure, 439

Musso, R. E. (see also Hinton, D. M.), 53
Myosin crossbridge lattice structure in insect flight muscle, 123
— gene duplication repeats, 15
— and rod minifilaments, crossbridge release and α-helix–coil transition, 455
—, scallop, electron microscopy; location of regulatory light chains, 723
—–actin interactions visualized by the quick-freeze, deep-etch replica technique, 97

Nisonoff, A. (see also Amit, A. G.), 637
Nuclear magnetic resonance measurements of intramolecular proton-proton distance constraints, and use of pseudo-structures for the 20 common amino acids in studies of protein conformation, 949

Ohlendorf, D. H., Anderson, W. F., Lewis, M., Pabo, C. O. & Matthews, B. W., Comparison of the structures of Cro and λ repressor proteins from bacteriophage λ, 757
Orlova, E. V. (see also Kiselev, N. A.), 345

Pabo, C. O. (see also Ohlendorf, D. H.), 757
Palmer, R. A. (see also Borkakoti, N.), 743
Pancreatic ribonuclease-A specificity, an X-ray study of a protein–nucleotide complex, 743
Partition of unit-copy miniplasmids to daughter cells. P1 and F miniplasmids contain discrete, interchangeable sequences sufficient to promote equipartition, 353
— — -— — — — —. The partition region of miniplasmid P1 encodes an essential protein and a centromere-like site at which it acts, 373
Pederson, T. (see also Roberts, M. P.), 975
Peeters, B. P. H., Konings, R. N. H. & Schoenmakers, J. G. G., Characterization of the DNA binding protein encoded by the N specific filamentous *Escherichia coli* phage IKe. Binding properties of the protein and nucleotide sequence of the gene, 197
Peptide NH protons, most slowly exchanging, in residues 1 to 19 of ribonuclease S, nature and locations, 281
Phage IKe, *Escherichia coli*, characterization of the encoded DNA binding protein, 197
Pirrotta, V. (see also Spena, A.), 799
Plastocyanin, poplar, oxidized, structure at 1·6 Å resolution, 521
Platt, T. (see also Horowitz, H.), 775
Poljak, R. J. (see also Amit, A. G.), 637
Poplar plastocyanin, oxidized, structure at 1·6 Å resolution, 521
Potassium hyaluronate, crystal structure of a tetragonal form containing extended chains, 813
— —, — — — tetragonal and orthorhombic forms containing sinuous chains, 829
— and sodium salts of chondroitin 4-sulfate, comparison of structures, 903
Proline content of ribonucleases does not affect their refolding, 619
Protein conformations by n.m.r. measurements of intramolecular proton–proton distance constraints, use of pseudo-structures for the 20 common amino acids, 949
—, DNA binding, encoded by the N-specific filamentous *Escherichia coli* phage IKe, characterization, 197
—, — —, gene *5*, from bacteriophage fd; refined structure, 565
—, membrane, a genetic approach to defining sites of interaction with different external agents, 507
—, T4 DNA helix destabilizing, (gp32*I), crystal, structural analysis by electron microscopy, 235
—–nucleotide complex, an X-ray study; specificity of pancreatic ribonuclease-A, 743
Protons, amide, H-bonded, in the 3 to 13 helix of ribonuclease S, exchange behaviour, 299
—, NH, peptide, most slowly exchanging, in residues 1 to 19 of ribonuclease S, nature and locations, 281
Proton–proton intramolecular distance constraint measurement by n.m.r., and use of pseudo-structures for the 20 common amino acids in studies of protein conformations, 949
Pseudo-structures for the 20 common amino acids for use in studies of protein conformations by measurement of intramolecular proton–proton distance constraints with n.m.r., 949
Pustovskikh, A. I. (see also Kiselev, N. A.), 345

Quick-freeze, deep-etch replica technique for visualization of actin–myosin interactions, 97

Raghunathan, S. (see also Arnott, S.), 861
— (see also Mitra, A. K.), 829
Reisler, E. (see also Applegate, D.), 455
Repeats, gene duplication, in the myosin rod, 15

Repressor proteins, cro and λ, from bacteriophage λ, comparison of structures, 757
RNA genes, small nuclear, U2, *Drosophila melanogaster*, isolation and characterization, 691
—, 5 S ribosomal, open and closed, the only two universal structures encoded in the nucleotide sequences, 1
—, ribosomal, 16 S, isolated, and its complexes with ribosomal proteins, shape and compactness, 409
—, 4·5 S, and the B1 family of repeated DNA in the 5′ flanking region of the mouse β-globin gene, a related sequence, 975
RNAs, ribosomal, small subunit, higher order structure in the 3′-minor domain, 249
Ribosomal 5 S RNA, open and closed, the only two universal structures encoded in the nucleotide sequences, 1
— 16 S RNA, isolated, and its complexes with ribosomal proteins, shape and compactness, 409
— 50 S subunit, computer averaging of electron micrographs, 345
— small subunit RNAs, higher order structure in the 3′-minor domain, 249
Ribonuclease refolding is independent of sequence variations, proline content and glycosylation, 619
—-A, pancreatic, specificity; an X-ray study of a protein–nucleotide complex, 743
— S, exchange behaviour of the H-bonded amide protons in the 3 to 13 helix, 299
— —, nature and location of the most slowly exchanging peptide NH protons in residues 1 to 19, 281
Roberts, M. P., Falvey, E., Calvet, J. P. & Pederson, T., A sequence related to 4·5 S RNA and the B1 family of repeated DNA in the 5′ flanking region of the mouse β-globin gene, 975
Rogero, J. R. (see also Englander, J. J.), 325

Salmonella typhimurium, nucleotide sequence of the *trpD* and *trpC* genes, 775
Saludjian, P. (see also Amit, A. G.), 637
Scallop myosin, electron microscopy; location of regulatory light chains, 723
Schmid, F. X. (see also Krebs, H.), 619
Schoenmakers, J. G. G. (see also Peeters, B. P. H.), 197
Schollmeier, K. (see also Hillen, W.), 707
Schon, E. A. (see also Shapiro, S. G.), 31
Sedelnikova, S. E. (see also Serdyuk, I. N.), 409
Selivanova, O. M. (see also Kiselev, N. A.), 345
Sequence and linkage of the goat ϵ^{I} and ϵ^{II} β-globin genes, 31
—, nucleotide, of the gene specifying the IKe DNA binding protein, 197
—, —, — — *trpD* and *trpC* genes of *Salmonella typhimurium*, 775
— related to 4·5 S RNA and the B1 family of repeated DNA in the 5′ flanking region of the mouse β-globin gene, 975
— variations of ribonucleases do not affect refolding, 619
Sequences, DNA, of yeast H3 and H4 histone genes from two non-allelic gene sets encode identical H3 and H4 proteins, 663
—, nucleotide, open and closed 5 S ribosomal RNA, the only two universal encoded structures, 1
—, zein, genomic, adjacent: structure, organization and tissue-specific restriction pattern, 799
Serdyuk, I. N., Agalarov, S. Ch., Sedelnikova, S. E., Spirin, A. S. & May, R. P., Shape and compactness of the isolated ribosomal 16 S RNA and its complexes with ribosomal proteins, 409
Shape and compactness of the isolated ribosomal 16 S RNA and its complexes with ribosomal proteins, 409
Shapiro, S. G., Schon, E. A., Townes, T. M. & Lingrel, J. B., Sequence and linkage of the goat ϵ^{I} and ϵ^{II} β-globin genes, 31
Sheehan, J. K. (see also Mitra, A. K.), 813, 829
Simmons, R. M. (see also Huxley, H. E.), 469
Skehel, J. J. (see also Wrigley, N. G.), 771
Skuratovskii, I. Ya. (see also Bartenev, V. N.), 217
Smith, M. M. & Andrésson, O. S., DNA sequences of yeast H3 and H4 histone genes from two non-allelic gene sets encode identical H3 and H4 proteins, 663
Smith, M. M. & Murray, K., Yeast H3 and H4 histone messenger RNAs are transcribed from two non-allelic gene sets, 641
Sodium and potassium salts of chondroitin 4-sulfate, comparison of structures, 903
Solano-Muñoz, F. (see also Bardsley, W. G.), 597
Spena, A., Viotti, A. & Pirrotta, V., Two adjacent genomic zein sequences: structure, organization and tissue-specific restriction pattern, 799
Spierer, P. (see also Hall, L. M. C.), 83
Spiess, E. (see also Alonso, A.), 691
Spirin, A. S. (see also Serdyuk, I. N.), 409

Squire, J. M. (see also Freundlich, A.), 439
Stel'maschuk, V. Ya. (see also Kiselev, N. A.), 345
Structural analysis of T4 DNA helix destabilizing protein (gp32*I) crystal by electron microscopy, 235
— implications of changes in the X-ray reflections from contracting muscle during rapid mechanical transients, 469
Structure of the *B* DNA cationic shell as revealed by an X-ray diffraction study of CsDNA; sequence-specific cationic stabilization of *B* form DNA, 217
—, higher order, in the 3′-minor domain of small subunit ribosomal RNAs from a gram negative bacterium, a gram positive bacterium and a eukaryote, 249
— of the myosin crossbridge lattice in insect flight muscle, 123
—, organization and tissue-specific restriction pattern for two adjacent genomic zein sequences, 799
— of oxidized poplar plastocyanin at 1·6 Å resolution, 521
—, refined, of the gene *5* DNA binding protein from bacteriophage fd, 565
—, three-dimensional, of the insect (*Lethocerus*) flight muscle M-band, 439
Structures of cro and λ repressor proteins from bacteriophage λ, a comparison, 757
— — the potassium and sodium salts of chondroitin 4-sulfate, a comparison, 903
—, universal, encoded in the nucleotide sequences, open and closed 5 S ribosomal RNA, 1
Substrate inhibition of enzyme-catalysed reactions, a theoretical and Monte Carlo study of turning points in $v(S)$ graphs, 597
Symmetry, molecular, of glyceraldehyde-3-phosphate dehydrogenase from *Bacillus coagulans*, 963

T4 (see also Bacteriophage), 427
— DNA helix destabilizing protein (gp32*I) crystal, structural analysis by electron microscopy, 235
Tail fiber attachment in bacteriophage T4 assembly; identification of the baseplate protein to which they attach, 427
Tedesco, P. (see also Urig, M. A.), 427
Tetracycline resistance genes, Tn*10*-encoded, control of expression; equilibrium and kinetic investigation of the regulatory reactions, 707
3′-minor domain of small subunit ribosomal RNAs, higher order structure, 249
Tn*10*-encoded tetracycline resistance genes, control of expression; equilibrium and kinetic investigation of the regulatory reactions, 707
Townes, T. M. (see also Shapiro, S. G.), 31
Transcription, *in vitro*, specific, of the insertion sequence IS2, 53
Transcripts, genes and bands in 315,000 base-pairs of *Drosophila* DNA, 83
Trifonov, E. N. & Bolshoi, G., Open and closed 5 S ribosomal RNA, the only two universal structures encoded in the nucleotide sequences, 1
trpD and *trpC* genes of *Salmonella typhimurium*, nucleotide sequences, 775

U2 small nuclear RNA genes, *Drosophila melanogaster*, isolation and characterization, 691
Unfolding unit, allosterically sensitive, in hemoglobin, identification, 325
Universal structures encoded in the nucleotide sequences, open and closed 5 S ribosomal RNA, 1
Urig, M. A., Brown, S. M., Tedesco, P. & Wood, W. B., Attachment of tail fibers in bacteriophage T4 assembly. Identification of the baseplate protein to which tail fibers attach, 427

Van Arsdell, J. (see also Horowitz, H.), 775
Vasiliev, V. D. (see also Kiselev, N. A.), 345
Vibert, P. (see also Flicker, P. F.), 723
Viotti, A. (see also Spena, A.), 799
Volkova, L. I. (see also Bartenev, V. N.), 217
$v(S)$ graphs, a theoretical and Monte Carlo study of turning points; inhibition of enzyme-catalysed reactions by excess substrate, 597

Wallimann, T. (see also Flicker, P. F.), 723
Wider, G. (see also Braun, W.), 921
Wood, W. B. (see also Urig, M. A.), 427
Wright, A. J. (see also Bardsley, W. G.), 597
Wrigley, N. G., Brown, E. B. & Skehel, J. J., Electron microscopic evidence for the axial rotation and inter-domain flexibility of the Fab regions of immunoglobulin G, 771
Wüthrich, K., Billeter, M. & Braun, W., Pseudo-structures for the 20 common amino acids for use in studies of protein conformations by measurements of intramolecular proton–proton distance constraints with nuclear magnetic resonance, 949
— (see also Braun, W.), 921

X-ray diffraction of CsDNA reveals the structure of *B* DNA cationic shell; sequence-specific cationic stabilization of *B* form DNA, 217
— reflections from contracting muscle, changes during rapid mechanical transients, and their structural implications, 469
— study of a protein–nucleotide complex; specificity of pancreatic ribonuclease-A, 743

Yeast H3 and H4 histone genes, DNA sequences from two non-allelic gene sets encode identical H3 and H4 proteins, 663
— — — — — messenger RNAs are transcribed from two non-allelic gene sets, 641

Zein sequences, genomic, adjacent; structure, organization and tissue-specific restriction patterns, 799